

QMW-92-7

May 1992

SINGULAR CONTRACTIONS OF W -ALGEBRAS

C. M. HULL AND L. PALACIOS^{*}

*Physics Department, Queen Mary and Westfield College,
Mile End Road, London E1 4NS, United Kingdom.*

ABSTRACT

Many W -algebras (e.g. the W_N algebras) are consistent for all values of the central charge except for a discrete set of exceptional values. We show that such algebras can be contracted to new consistent degenerate algebras at these exceptional values of the central charge.

^{*} supported by CONACyT-MEXICO, grant No. 52361

Many W -algebras, such as the W_N algebras [1- 5] which are generated by currents of spin $2, 3, \dots, N$, have the property that they are defined for all values of the central charge, except for a discrete set of exceptional values. It is the purpose of this paper to investigate such W -algebras for these exceptional values of the central charge.

This work was originally stimulated by the paper [8] which claimed to give a construction of the W_4 algebra for $c = -2$, which appeared to contradict the results of [6, 7] who claimed that the W_4 algebra was defined and satisfied the Jacobi identities for all values of c except a discrete set of exceptional values, which unfortunately included the value $c = -2$. This suggested that the $c = -2$ algebra of [8] could not be the usual W_4 algebra, and raised the question as to what it might be. We will return to this algebra at the end of the paper.

The W_3 algebra is given by

$$\begin{aligned}
[L_n, L_m] &= \frac{c}{2} \frac{n(n^2 - 1)}{6} \delta_{n+m} + (n - m) L_{n+m} \\
[L_n, W_m] &= (2n - m) W_{n+m} \\
[W_n, W_m] &= \frac{c}{3} \frac{(n^3 - n)(n^2 - 4)}{5!} \delta_{n+m} \\
&\quad + \frac{(n - m)}{30} [2(n + m)^2 - 5nm - 8] L_{n+m} \\
&\quad + \frac{16}{22 + 5c} (n - m) \Lambda_{n+m}
\end{aligned} \tag{1}$$

where

$$\Lambda_m = \sum_n : L_{m-n} L_n : - \frac{3}{10} (m + 3)(m + 2) L_m, \tag{2}$$

For any currents A, B with modes A_n, B_n , we define the usual normal ordered product $: A_n B_m :$ by [1, 3].

$$: A_n B_m := \begin{cases} A_n B_m, & \text{if } m \geq n \\ B_m A_n, & \text{if } m < n \end{cases} \tag{3}$$

The form of the algebra is uniquely determined by requiring that the Jacobi identity be satisfied and that the central charge terms have a conventional normalisation

[1]. Note, however, that the structure constants become divergent if $c = \frac{-22}{5}$, unless the spin-four quasi-primary field Λ given by (2) vanishes or can be set to zero. However, the fact that the form of the coupling constants is fixed by requiring a conventional normalisation of the currents suggests that it might be possible to absorb the divergence occurring for $c = \frac{-22}{5}$ into a rescaling of the generators. To this end, we consider the redefinition

$$W_n \rightarrow W'_n = \sqrt{5c + 22} W_n \quad (4)$$

The new generator W'_n is still a spin-three primary field, but the $[W', W']$ commutator is now

$$\begin{aligned} [W'_n, W'_m] = & \frac{c(22 + 5c)}{3} \frac{(n^3 - n)(n^2 - 4)}{5!} \delta_{n+m} \\ & + \frac{(n - m)(22 + 5c)}{30} [2(n + m)^2 - 5nm - 8] L_{n+m} \\ & + 16(n - m) \Lambda_{n+m} \end{aligned} \quad (5)$$

For all values of c not equal to $-22/5$, this gives an algebra which is equivalent to the original W_3 algebra. However, for the rescaled algebra, it is now possible to take the limit $c \rightarrow \frac{-22}{5}$ while keeping W' finite and obtain a non-singular algebra. The resulting algebra has a primary spin-three generator W' satisfying

$$[W'_n, W'_m] = 16(n - m) \Lambda_{n+m} \quad (6)$$

The absence of any central charge term in this commutator implies that W' is null, in the sense that the vacuum expectation value $\langle W'(z)W'(w) \rangle$ vanishes and $W'_{-n} | \phi \rangle$ is a zero-norm state for any state $| \phi \rangle$ satisfying the condition $W'_n | \phi \rangle = 0$ for $n > 0$. The fact that the algebra generated by L_n, W'_n satisfies the Jacobi identities for all $c \neq -22/5$ implies, by continuity, that the new algebra obtained in the $c \rightarrow \frac{-22}{5}$ limit also satisfies the Jacobi identities. The construction of this new algebra as a limit of rescaled algebras is similar to a Wigner-Inonu contraction.

In general, for a W -algebra with generators W_n^A of conformal dimension Δ_A (labelled by $A = 1, 2, \dots$), the algebra will take the form

$$[W_m^A, W_n^B] = g^{AB} m(m^2 - 1) \dots (m^2 - (\Delta_A - 1)^2) \delta_{m+n} + \dots \quad (7)$$

(plus W^A -dependent terms) for some constant matrix g^{AB} , so that $\langle W^A W^B \rangle \propto g^{AB}$, and we shall refer to g^{AB} as the metric of the algebra. On the right hand side of (7), terms involving the generators have been suppressed. For algebras with unitary representations g^{AB} should be positive definite and should vanish if $\Delta_A \neq \Delta_B$. Then a basis can be chosen to diagonalize g^{AB} and it is conventional to normalise the generators so that

$$[W_m^A, W_n^B] = \frac{c}{\Delta(2\Delta - 1)!} \delta^{AB} \delta_{m+n} m(m^2 - 1) \dots (m^2 - (\Delta - 1)^2) + \dots$$

for some central charge c . Given this normalisation, requiring the Jacobi identities to hold fixes the form of most of the known W -algebras uniquely. In many cases, the resulting structure constants are singular for certain discrete values of the central charge c and in these cases it is again possible to find a contracted algebra which is non-singular at these special values of the central charge. In each case, the contracted algebra has a metric g^{AB} which has zero eigenvalues and hence includes null generators.

The W_4 algebra is given by [6]

$$\begin{aligned} [W_n^A, W_m^B] = & \frac{c}{\Delta} \frac{m(m^2 - 1) \dots (m^2 - (\Delta - 1)^2)}{(2\Delta - 1)!} \delta^{AB} \delta_{m+n} \\ & + f^{AB}{}_C p_{\Delta_A \Delta_B \Delta_C}(n, m) W_{n+m}^C \\ & + C^{AB}{}_a p_{\Delta_A \Delta_B \Delta_a}(n, m) \Lambda_{n+m}^a \end{aligned} \quad (8)$$

where W_n^A ($W_n^2 \equiv L_n, W_n^3 \equiv W_n, W_n^4 \equiv V_n$) are the modes of the primary fields of spin- Δ_A , ($\Delta_A = 2, 3$ and 4 , respectively). The $p_{\Delta_A \Delta_B \Delta_C}(n, m)$ are certain ‘universal’ polynomials and the Λ^a ($a = 1, \dots, 7$) are composite operators constructed

from the primary fields, given by

$$\begin{aligned}
\Lambda^1 = \Lambda &=: W^2 W^2 : -\frac{3}{10} \partial^2 W^2 \\
\Lambda^2 = A &=: W^3 W^2 : -\frac{3}{14} \partial^2 W^3 \\
\Lambda^3 = \Gamma &=: W^2 \partial^2 W^2 : -\partial : W^2 \partial W^2 : +\frac{2}{9} \partial^2 : W^2 W^2 : -\frac{1}{42} \partial^4 W^2 \\
\Lambda^4 = \Delta &=: \Lambda W^2 : -\frac{1}{6} \partial^2 \Lambda \\
\Lambda^5 = \Omega &=: W^4 W^2 : -\frac{1}{6} \partial^2 W^4 \\
\Lambda^6 = \Pi &=: W^3 W^3 : -\frac{1}{84} \partial^4 W^2 - \frac{40}{9(5c+22)} \partial^2 \Lambda - \frac{5}{36} f_4^{33} \partial^2 W^4 \\
\Lambda^7 = \Theta &=: W^3 \partial W^2 : -\frac{2}{5} \partial : W^3 W^2 : +\frac{1}{20} \partial^3 W^3
\end{aligned} \tag{9}$$

with $:AB:$ indicating the normal ordering of the operators A and B expressed by (3) when they are decomposed in terms of their modes A_n and B_n .

The structure constants f^{AB}_C and C^{AB}_c that depend on the central charge [6] are given below:

$$\begin{aligned}
(f_4^{33})^2 &= \frac{16(7c+114)(c+2)}{3(5c+22)(c+7)} \\
(f_4^{44})^2 &= \frac{27(c^2+c+218)^2}{(7c+114)(5c+22)(c+7)(c+2)} \\
f_3^{34} &= \frac{3}{4} f_4^{33}
\end{aligned} \tag{10}$$

$$\begin{aligned}
C_{\Lambda}^{33} &= \frac{32}{5c+22} \\
C_{\Lambda}^{44} &= \frac{42}{5c+22} \\
C_{\Gamma}^{44} &= \frac{3(19c-582)}{20(7c+114)(c+2)} \\
C_{\Delta}^{44} &= \frac{96(9c-2)}{(7c+114)(5c+22)(c+2)} \\
C_{\Omega}^{44} &= \frac{36(c^2+c+218)}{(5c+22)(c+2)(c+7)} (f_4^{44})^{-1} \\
C_{\Pi}^{44} &= \frac{45(5c+22)}{2(7c+114)(c+2)} \\
C_A^{34} &= \frac{39}{7c+114} f_4^{33}; \quad C_{\Theta}^{34} = \frac{3}{4(c+2)} f_4^{33}
\end{aligned} \tag{11}$$

The algebra satisfies the Jacobi identities for all values of c , except for the isolated points $c = -2, -\frac{22}{5}, -7, -\frac{114}{7}$ [6] (although in [7] it is claimed that the Jacobi identities are satisfied for all values of the central charge except $c = -2, -\frac{22}{5}, -7, -\frac{114}{7}, \frac{1}{2}, -\frac{68}{7}, -\frac{2}{5}$). The structure constants become complex in the intervals $-114/7 \leq c \leq -7$ and $-22/5 \leq c \leq -2$.

Although the W_4 algebra does not exist at these exceptional values of the central charge, we can again define new contracted algebras at these values of c . Consider first rescaling the spin-4 generator, V_n , by $V_n \rightarrow \sqrt{5}\sqrt{c+2}V_n$ and then taking the limit $c \rightarrow -2$. This gives a contracted algebra generated by the Virasoro generators and the spin three and four primary fields W, V satisfying

$$\begin{aligned}
[W_n, W_m] &= \frac{-2}{3} \frac{n(n^2-1)(n^2-4)}{5!} \delta_{n+m} \\
&\quad + \frac{(n-m)}{30} [2(n+m)^2 - 5nm - 8] L_{n+m} \\
&\quad + \frac{4}{3} (n-m) \Lambda_{n+m} + \frac{10}{3} (n-m) V_{n+m} \\
[V_n, V_m] &= \frac{33}{180} (n-m) [(n-m)^2 + nm - 7] V_{n+m} \\
&\quad + \frac{(n-m)}{50} \left[-\frac{93}{20} \Gamma_{n+m} - 8\Delta_{n+m} + 100\Omega_{n+m} + \frac{27}{2} \tilde{\Pi}_{n+m} \right] \\
[W_n, V_m] &= \Theta_{n+m}
\end{aligned} \tag{12}$$

where $\Gamma_n, \Delta_n, \Omega_n$, and Θ_n are given by (9), and $\tilde{\Pi} =: WW : -\frac{1}{84}\partial^4 L - \frac{10}{27}\partial^2 \Lambda - \frac{25}{36}\partial^2 V$.

Similarly, it is possible to obtain a contracted W_4 algebra for $c = -114/7$. However, after the contractions has been done, some of the new structure constants in the commutators of the spin three and four primary fields are not real numbers. After the transformation

$$W \rightarrow -i\sqrt{\frac{14}{13}}W \quad (13)$$

$$V \rightarrow 4i\sqrt{\frac{42}{5}}V, \quad (14)$$

the structure constant are all real and the contracted W_4 algebra at $c = -\frac{114}{7}$ is given by the following set of commutators (in addition to the Virasoro algebra and the ones that state the fact that the generators W and V are primary fields)

$$\begin{aligned} [W_m, W_n] &= p_{334}(m, n)[4V_{m+n} + \frac{1}{2}\Lambda_{m+n}] \\ &\quad - \frac{13}{7}p_{332}(m, n)L_{m+n} \\ &\quad + \frac{247}{49}\binom{m+2}{5}\delta_{m+n} \end{aligned} \quad (15)$$

$$[W_m, V_n] = \frac{5}{4}p_{345}(m, n)\Lambda_{m+n} \quad (16)$$

$$\begin{aligned} [V_m, V_n] &= p_{446}(m, n)[\frac{3}{4}\tilde{\Pi}_{m+n} + \frac{1}{8}\Delta_{m+n} - \frac{39}{80}\Gamma_{m+n}] \\ &\quad - p_{444}(m, n)\frac{50}{3} \cdot \frac{275}{7}V_{m+n} \end{aligned} \quad (17)$$

where

$$\tilde{\Pi} =: WW : -\frac{5}{9}\partial^2 V - \frac{5}{72}\partial^2 \Lambda + \frac{13}{1176}\partial^4 L.$$

After the rescaling $W \rightarrow (c+7)^{-q}W$ and $V \rightarrow (\sqrt{3}(c+7))^{-p}V$, to obtain a well defined limit of the structure constants at $c = -7$, the exponents p and q must

satisfy the following relations

$$\begin{aligned} p = q \quad \text{with} \quad q \geq \frac{1}{2} \\ \text{or} \quad p = 2q - \frac{1}{2}; \quad q > \frac{1}{2} \end{aligned} \tag{18}$$

The first relation with $q = p = \frac{1}{2}$ gives rise to the following commutation relations

$$\begin{aligned} [W_n, W_m] &= \frac{10}{3}(n-m)V_{n+m} \\ [V_n, V_m] &= \frac{n-m}{3}[(n-m)^2 + nm - 7]V_{n+m} \\ &\quad + \frac{n-m}{2}[4\Omega_{n+m} + \frac{9}{10}\tilde{\Pi}_{n+m}] \\ [W_n, V_m] &= \frac{15}{84}(5n^3 - m^3 - 5mn^2 + 3m^2n - 17n + 9m)W_{n+m} \\ &\quad + \frac{12}{5}(3n + 2m)A_{n+m} - 3\Theta_{n+m} \end{aligned} \tag{19}$$

where $\tilde{\Pi} =: WW : -\frac{25}{27}\partial^2 V$. If $p = q > \frac{1}{2}$ then the contracted commutation relations are

$$\begin{aligned} [W_n, W_m] &= 0 = [W_n, V_m] \\ [V_n, V_m] &= \frac{9}{20}(n-m)\tilde{\Pi}_{n+m} \end{aligned} \tag{20}$$

where $\tilde{\Pi} =: WW :$.

The second relation in (18) allows the following limit

$$\begin{aligned} [V_n, V_m] &= 0 = [W_n, V_m] \\ [W_n, W_m] &= \frac{10}{3}(n-m)V_{n+m}. \end{aligned} \tag{21}$$

The contraction for the case $c = -\frac{22}{5}$ can be obtained through the rescaling $W \rightarrow (5c - 22)^{-q/2}W$ and $V \rightarrow (5c - 22)^{-p/2}V$ with $(p, q) \in \{(p, q)/p \geq \frac{1}{2}, q \geq \frac{1}{2}, p \geq q - \frac{1}{2} \text{ and } p \leq 2q - \frac{1}{2}\}$. The result is a set of null field commutation relations for the generators of conformal dimension three and four, with some of the coefficients complex.

To summarise, we have seen that W -algebras that are consistent for all but a discrete set of exceptional values of the central charge can be contracted to algebras that can be continued consistently to these values of c , resulting in new algebras at these c -values. These new algebras have a degenerate metric, so that some of the generators have become null and the rank of the algebra has been reduced.

We now return to the paper [8], in which W_N algebras for $c = -2$ are claimed to have been constructed. For the $N = 4$ case, for example, quasi-primary currents T, W, V of spins 2, 3 and 4 are constructed from free fermions and it is claimed that they satisfy a closed algebra which also satisfies the Jacobi identities and contains the Virasoro algebra with $c = -2$. It is usually assumed in such cases that each of the quasi-primary currents of spins greater than 2 can be modified by expressions involving composites constructed from the currents and their derivatives in such a way as to obtain primary currents. The algebra in this primary basis should then again be closed and satisfy the Jacobi identities. If this were the case, and if all the currents were non-null, then the algebra should be the standard W_4 algebra given by (8) for some value of the central charge. However, the Virasoro generators are unmodified and so still satisfy the Virasoro algebra with $c = -2$, implying that the whole W_4 algebra would have $c = -2$. However, as we have seen, the standard W_4 algebra does not exist at $c = -2$, so it follows that at least one of our assumptions must be wrong. The quasi-primary fields in the original algebra of [8] were all non-null, *i.e.* the expectation values $\langle TT \rangle$, $\langle WW \rangle$ and $\langle VV \rangle$ were all non-zero. This suggests that the problem must come from the change of basis from quasi-primary to primary currents, and that some of the primary currents become null. We now show that this is indeed the case. The spin-three current W of [8] is in fact primary and needs no modification, but the spin-four current V is not primary. It is straightforward to show that the current

$$V' = V - :TT: - \frac{3}{10}\partial^2 T \quad (22)$$

is in fact primary, so that a primary basis for the algebra is given by T, W, V' . However, for the fermionic realisation of [8], the current V' vanishes identically,

so that the algebra collapses to the W_3 algebra at $c = -2$, which is of course a special case of the contracted W_4 algebra at $c = -2$ constructed above (and it is possible formally to re-introduce a null primary spin-four current V' to obtain precisely this contracted algebra). Similarly, for each of the ' W_N ' algebras of [8] with $N > 3$, there is again the problem that none of the W_N algebras with $N > 3$ exist for $c = -2$. The resolution is again that, on changing to a basis of primary currents, each of the currents of spin greater than 3 vanish identically, so that for each case the algebra that is constructed is just W_3 (or a singular contraction of W_N) and so the problem that is avoided.

After this work was completed, we learned of related work by Niedermaier [9], in which he considers W_N algebras in a basis of quasi-primary currents, and shows that these algebras are consistent for all values of c , and in particular the structure constants are non-singular functions of c . However, the rank of the algebra (essentially the number of non-null generators) which is $N - 1$ for generic values of c , becomes smaller for special values of c , e.g. for W_4 , the rank of the algebra becomes less than three for $c = 0, -2, -7, -114/7$. Further, the change of basis from quasi-primary to primary currents which is possible for generic values of c , becomes singular at $c = -2, -22/5, -7, -114/7$. By relaxing the conventional normalisation condition on the primary fields, it is possible to obtain a non-singular change of basis to obtain one of the contracted algebras described above for $c = -2, -7, -114/7$ but this is not possible for the value $c = -22/5$. Thus the W_4 algebra of [9] is equivalent to the standard W_4 algebra for non-exceptional values of the central charge but is equivalent to the contracted W_4 algebras described above.

Acknowledgments We would like to thank Horst Kausch and Chris Pope for discussions and one of us (LP) would like to thank CONACYT-México for financial support.

REFERENCES

1. A.B. Zamolodchikov, Teor. Mat. Fiz. **65** (1985) 1205.
2. V.A. Fateev and S. Lykanov, Intern. J. Mod. Phys. **A3** (1988) 507.
3. F. Bais, P. Bouwknegt, M. Surridge and K. Schoutens, Nucl. Phys. **B304** (1988) 348, 371.
4. A. Bilal and J.-L. Gervais, Phys. Lett. **206B** (1988) 412; Nucl. Phys. **B314** (1989) 646; Nucl. Phys. **B318** (1989) 579; Ecole Normale preprint LPTENS 88/34.
5. C.M. Hull, Nucl. Phys. **353 B** (1991) 707.
6. R. Blumenhagen, M. Flohr, A. Kliem, W. Nahm, A. Recknagel and R. Varnhagen, Bonn University preprint BONN-HE-90-05(1990).
7. H. G. Kausch and G. M. T. Watts, Nucl. Phys. **B354** (1991) 740.
8. H. Lu, C. N. Pope X. Shen and X.J. Wang, Phys. Lett. **B267**(1991) 356 - 361.
9. M.R. Niedermaier, DESY preprint 91-148.